

NewportCare Medical Group

Lucas Wymore MD


Sports Medicine

Office Phone: 949-491-9991

40700 California Oaks Drive, Suite 205

Murrieta, CA 92592

Fax: 949-612-9795



NewportCare®

BICEPS TENDON ALLOGRAFT RECONSTRUCTION REHABILITATION PROTOCOL

	RANGE OF MOTION	IMMOBILIZER	THERAPEUTIC EXERCISE
PHASE I 0 - 3 weeks	Keep brace locked at 60 flexion	Locked in neutral - worn at all times	Gentle wrist and shoulder ROM
PHASE II 3 - 6 weeks	Advance extension in brace gradually	Worn at all times (including exercise) - removed for hygiene	Continue with wrist and shoulder ROM, begin active extension to full, NO active flexion, gentle joint mobilizations
PHASE III 6 - 9 weeks	Active extension to 0 in brace	Worn at all times (including exercise) - removed for hygiene	Continue to maintain wrist and shoulder flexibility, begin rotator cuff/deltoid isometrics, progress active extension in brace
PHASE IV 9 - 12 weeks	Gently advance ROM to tolerance	None	Begin active flexion and extension against gravity, advance strengthening in phase III to resistive, maintain flexibility/ROM
PHASE V 12 weeks - 6 months	Gradual return to full and pain-free	None	Begin gentle flexion strengthening, advance activities in phase IV
PHASE VI 6 months and beyond	Full and pain-free	None	Return to full activity