



**Lucas Wymore, MD**  
Sports Medicine  
40700 California Oaks Drive Suite 205  
Murrieta CA 92562

Office Phone: 949-491-9991  
Office Fax: 949-612-9795  
Email: [Wymore@newportcare.org](mailto:Wymore@newportcare.org)

## **PATIENT GUIDE TO SHOULDER INSTABILITY**

### **What is shoulder instability?**

Shoulder instability is a cause of shoulder pain. The shoulder can dislocate (the ball comes completely out of the socket) or subluxate (the ball comes partially out of the socket). After this happens, pain and inability to use the shoulder occurs. Shoulder instability is the result of damage to the shoulder ligaments that keep the ball connected to the socket.

### **How does shoulder instability occur?**

Many times, the shoulder ligaments tear from a single violent injury, such as a football tackle or motor vehicle accident. However, they can also be stretched from repeated strenuous use, such as with swimming or throwing.

### **How do I know my shoulder is unstable?**

In many cases, the patient knows they have shoulder instability because they need someone to reduce the joint back into place (such as an emergency room physician). In other cases, the patient primarily complains of pain in the shoulder, with subtle feelings of slipping. The examination in the office usually confirms the diagnosis of shoulder instability.

### **Do I need x-rays, MRI's or any other test?**

A set of x-rays is usually ordered to make sure there are no fractures (broken bone) in the shoulder. Commonly, a small fracture in the area of the dislocation can occur on the humeral head, called a Hill-Sachs lesion. A MRI can sometimes be helpful to confirm the tear of the ligament off of the bone (Bankart lesion), and evaluate for other shoulder problems. Usually, the MRI is performed with dye (gadolinium) to help see the ligaments.

### **Is there other damage to the shoulder in cases of instability?**

The labrum (where the shoulder ligaments attach) is usually torn in cases of shoulder instability. In addition, there are sometimes fractures of the ball or socket of the shoulder. Rarely, there is also a rotator cuff tear in cases of instability. A rotator cuff tear is more common in patients over the age of 40 with a shoulder dislocation.

### **What are the treatment options for shoulder instability?**

The treatment primarily depends on the patient's age, activity level, and number of dislocations of the shoulder. When a patient dislocates their shoulder for the first time, they can commonly be treated in a sling with early rehabilitation. However, for patients with repeat episodes of instability, the usual treatment is surgery to fix the ligaments.

### **How is shoulder instability treated with surgery?**

Shoulder instability is usually repaired with arthroscopic techniques. The arthroscope is a fiber optic instrument (narrower than a pen) which is put into the joint through small incisions. A camera is attached to the arthroscope and the image is viewed on a TV monitor. The arthroscope allows me to fully evaluate the entire shoulder joint, including the ligaments, the rotator cuff, and the cartilage surface. Small instruments ranging from 3-5 millimeters in size are inserted through additional incisions so that I can feel the joint structures for any damage, diagnose the injury, and then repair, reconstruct, or remove the damaged tissue.

In shoulder instability surgery, the damaged labrum and ligaments are identified and then repaired back to the socket. This is usually performed by using suture anchors (a screw with sutures attached) to sew the ligaments back in place.

Occasionally, if the damage is severe or longstanding, the surgery needs to be performed with an open incision. This is a larger incision on the front of the shoulder used to get to the shoulder joint and repair the ligaments.

### **What are some of the possible complications?**

While complications are not common, all surgery has associated risk. One possible complication includes stiffness of the shoulder after surgery. The use of arthroscopic techniques attempts to limit these complications. Other complications include recurrent instability, an infection, bleeding, nerve damage, or problems with the anesthesia.

### **What kind of anesthesia is used?**

I use a combination of general anesthesia and regional anesthesia. Before the surgery, the anesthesiologist will inject some novocaine around the nerves of the shoulder. This numbs the arm and helps to control your pain after surgery. In addition, you go to sleep (general anesthesia) to help keep you comfortable during surgery.

### **What do I need to do to prepare for surgery?**

Our staff will help to set up the surgery through your insurance company and will instruct you on any paperwork that may be necessary.

Prior to your surgery, you may be asked to get several medical tests, done on an outpatient basis. Most patients need some minor blood tests. If you are over age 45, you may require an EKG. Some patients need to see an internist or their family doctor to obtain clearance for surgery.

The night before the surgery, a member of our staff will contact you about what time to arrive for surgery. You may not eat or drink anything after midnight the night before your surgery.

### **How long will I be in the hospital?**

Almost all patients are able to have surgery and go home the same day. Occasionally, patients will be admitted for an overnight stay.

### **What happens the day of surgery?**

The morning of your surgery you will be admitted and taken to a pre-operative holding area where you are prepared for surgery.

You will be asked several times which extremity I am operating on. Please note that you are asked this question many times on purpose. I will initial the operative arm prior to surgery.

After the operation, you will be taken to the recovery room to be monitored. Once the effects of anesthesia have worn off and your pain is under good control, you can see your family and finish recovering. You will be given all of your post-operative instructions and pain medication before leaving.

Please be aware that the process of getting checked in, prepared for surgery, undergoing the operation, and recovering from anesthesia takes the majority of the day. I would recommend that you and your family members bring along some reading material to make the process easier for all.

### **How should I care for my shoulder after surgery?**

Prior to your discharge, you will be given specific instructions on how to care for your shoulder. In general, you can expect the following:

**Diet:** Resume your regular diet as soon as tolerated. It is best to start with clear liquids before advancing to solid food.

**Medication:** You will be given a prescription for pain medication.

**Bandage:** You will have a thick dressing on the shoulder. You will be instructed on when it can be removed, usually in 3 days.

**Showering:** You may shower after your dressing is removed, after 3 days. You cannot take a bath until the wounds are completely sealed, usually 2 – 3 weeks after surgery.

**Sling:** You will have a sling, which you will use for 4 - 6 weeks. You can remove it for grooming and physical therapy.

**Ice:** You may receive an ice machine that continually surrounds your shoulder with cold water. If not, you may apply ice over the dressings for 30 minutes every hour for several days. Do not use heat.

**Suture removal:** Your stitches will be removed at your office visit 7-14 days after surgery. Occasionally, sutures are used which resorb and do not need to be removed.

**Follow-up office visit:** You will be instructed on when to follow-up in the office. This is usually 7-14 days after surgery.

**Exercise:** You will be instructed on exercises you can do immediately after surgery. You will start physical therapy within 1 to 2 weeks after surgery.

**Return to work or school:** You can return to school or work within 3 – 5 days without using the affected arm. If you need the use of the arm to return, you may be out of work or school for a longer period of time.

### **What will rehabilitation involve?**

The rehabilitation is based on several goals: 1) allowing the tissue to heal; 2) regaining motion; 3) regaining strength; and 4) return to sports. The rehabilitation protocol for the physical therapist is attached for you to review.

### **When can I return to sports?**

In general, you will be allowed to return to sports in 6 months after surgery. You must have good motion, strength, and control of your shoulder and arm. How quickly you return to sports depends on several factors, including: 1) your own rate of healing; 2) the damage found at surgery; 3) if you have any complications; 4) how well you follow the post-operative instructions; 5) how hard you work in rehabilitation.

### **What is the success rate?**

Overall, the success rate for shoulder instability surgery ranges from 85 to 95% for attaining a stable shoulder.

## **Our commitment**

The entire NewportCare team is committed to you, the patient. We understand that this is a stressful time, and you may be anxious about your injury and the need for surgery. Please contact me with any questions about your injury or treatment plan.

### **Lucas Wymore, MD**

Office Phone: 949-491-9991

Office Fax: 949-612-9795

Email: [Wymore@newportcare.org](mailto:Wymore@newportcare.org)