number 596

may 2025

Our regular monthly meeting will be held at

Round Table Pizza

4200 Gosford Rd, Bakersfield, CA 93313

May 2

Social Hour @ 6:00p Meeting @ 7:00p

f

facebook.com/groups/syzygy



kernastro.org



kernastronomicalsociety@gmail.com

UPCOMING EVENTS

Star Party

May 24th at the Frazier Mountain Trailhead, weather permitting. Click for map

Club Events

May 3 - Annual Club Picnic at Jastro Park from 10a - 2p August 24 - Mount Wilson Observatory. **This event is full!**

NEW MEMBERS

Frank & Diana Ripepi and Bethany Banks have joined the Kern Astronomical Society! Please give them a warm welcome when you see them next!

EUROPA CLIPPER

Presented by Sarah Elizabeth McCandless

NASA's Europa Clipper spacecraft launched in October 2024 and will conduct a detailed survey of Jupiter's icy moon Europa to determine whether there are places below the surface that could support life. The spacecraft will arrive at Jupiter in 2030, and once in orbit around Jupiter, it will conduct nearly 50 flybys



of Europa, shifting its flight path for each encounter to soar over a different location so that it eventually scans almost the entire moon.

Sarah Elizabeth McCandless works for NASA's Jet Propulsion Laboratory (JPL) as a navigation engineer. She has flown eleven different missions exploring our solar system, including the Mars Perseverance rover, Artemis I, Psyche, and Europa Clipper. Working as an "interplanetary Google maps", she ensures we know a spacecraft's location and can precisely predict its trajectory to ensure successful orbit insertion about or landing on another planet or body. In 2024, she was selected for NASA's Human Exploration Research Analog (HERA) Campaign 7 Mission 3 and served as an "analog astronaut", simulating a mission to Mars for 45 days at NASA's Johnson Space Center. McCandless earned a Bachelor of Science in Aerospace Engineering and minor in French at the University of Kansas, and earned a Master of Science in Aerospace Engineering at the University of Texas at Austin, focusing on orbital mechanics. Outside of work, she enjoys running, camping, piloting Cessna 172s and traveling with friends and family.



KAS Information

Since 1956, the Kern Astronomical Society has promoted community awareness of current events in astronomy and provides a forum for sharing of knowledge and experiences among amateur astronomers. Annual membership is \$25.00 which also provides membership in the Amateur Astronomical League, access to their newsletter (Reflector Magazine), and participation in observational programs.


Star Parties & Outreach

The Kern Astronomical Society typically has two Club Star Parties each month depending on the weather. Our Club Parties are held on Saturdays nearest the New Moon. We also host Public Star Parties at various locations around town during April - October. These parties are held on Saturdays nearest the first quarter Moon. In addition, we also host Lunar, Solar, and Planetary viewing for public schools. Requests may be directed to our Star Party Coordinator.

Club Equipment

The Kern Astronomical Society has telescopes and accessories (listed below) available for loan to Club Members in good standing. Members are encouraged to borrow the different types of telescopes in stock (especially if you are considering purchasing one). Trying out different sizes and types of telescopes can help you make an informed decision about purchases. If you have a Club telescope in your possession, you will be expected to participate in at least one public star party.

- 6" f/6, 8" f/6, 10" f/5.6, 13" f/4.5 Dobsonian Telescopes
- Parks Jovian 90
- 3½" f/13 Maksukov-Cassegrain
- 4" f/15 Unitron Refractor
- 8" Solar Filter
- Assorted Eyepieces



About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars. They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy - A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT) - A time system used by astronomers. Also known as Greenwich Mean Time. USA Eastern Standard Time (for example, New York) is 5 hours behind UT.

Variable Star - A star that changes brightness over a period of time.

NORTHERN HEMISPHERE MAY

₩

M104

γ Virginis

Capella

Arcturus

Easily Seen with the Naked Eve

Procyon	CMi	• Greek name meaning "before the dog" - rises before Sirius (northern latitudes). Dist=11.4 ly.
δ Cephei	Сер	Cepheid prototype. Mag varies between 3.5 & 4.4 over 5.366 days. Mag 6 companion.
Deneb	Cyg	 Brightest star in Cygnus. One of the greatest known supergiants. Dist=1,400±200 ly.
Castor	Gem	 Multiple star system with 6 components. 3 stars visible in telescope. Dist=52 ly.
Pollux	Gem	 With Castor, the twin sons of Leda in classical mythology. Dist=34 ly.
lpha Herculis	Her	Semi-regular variable. Magnitude varies between 3.1 & 3.9 over 90 days. Mag 5.4 companion.
Regulus	Leo	 Brightest star in Leo. A blue-white star with at least 1 companion. Dist=77 ly.
Vega	Lyr	• The 5th brightest star in the sky. A blue-white star. Dist=25.0 ly.

• Orange, giant K star. Name means "bear watcher". Dist=36.7 ly.

• The 6th brightest star. Appears yellowish in color. Spectroscopic binary. Dist=42 ly.

Antares Red, supergiant star. Name means "rival of Mars". Dist=135.9 lv.

Polaris • The North Pole Star. A telescope reveals an unrelated mag 8 companion star. Dist=433 ly. Spica

• Latin name means "ear of wheat" and shown held in Virgo's left hand. Dist=250 ly.

Easily Seen with Binoculars

M44	Cnc	0	Praesepe or Beehive Cluster. Visible to the naked eye. Dist=590±20 ly.
M3	CVn	\oplus	Easy to find in binoculars. Might be glimpsed with the naked eye.
μ Cephei	Сер	•	Herschel's Garnet Star. One of the reddest stars. Mag 3.4 to 5.1 over 730 days.
Mel 111	Com	0	Coma Berenices. 80 mag 5-6 stars in 5 deg. Dist=283 ly. Age=400 million years.
χ Cygni	Cyg	•	Long period pulsating red giant. Magnitude varies between 3.3 & 14.2 over 407 days.
M39	Cyg	0	May be visible to the naked eye under good conditions. Dist=900 ly.
ν Draconis	Dra	•	Wide pair of white stars. One of the finest binocular pairs in the sky. Dist=100 ly.
M13	Her	0	Best globular in northern skies. Discovered by Halley in 1714. Dist=23,000 ly.
M92	Her	0	Fainter and smaller than M13. Use a telescope to resolve its stars.
R Hydrae	Hya	•	Long period variable. Mag varies between 3.0 & 11.0 over 390 days. Brilliant red.
ε Lyrae	Lyr	•	Famous Double Double. Binoculars show a double star. High power reveals each a double.
R Lyrae	Lyr	•	Semi-regular variable. Magnitude varies between 3.9 & 5.0 over 46.0 days.
M12	0ph	0	Close to the brighter M10. Dist=18,000 ly.
M10	0ph	0	3 degrees from the fainter M12. Both may be glimpsed in binoculars. Dist=14,000 ly.
TC / C C E	0 1		and the second second second

IC 4665 Large, scattered open cluster. Visible with binoculars. 6633 Scattered open cluster. Visible with binoculars.

Μ4 A close globular. May just be visible without optical aid. Dist=7,000 ly.

M5 Fine globular star cluster. Telescope will reveal individual stars. Dist=25,000 ly.

Mizar & Alcor Good eyesight or binoculars reveals 2 stars. Not a binary. Mizar has a mag 4 companion. Cr 399

Coathanger asterism or "Brocchi's Cluster". Not a true star cluster. Dist=218 to 1,140 ly.

Telescopic Objects							
ε Boötis	Boo	•	Red giant star (mag 2.5) with a blue-green mag 4.9 companion. Sep=2.8". Difficult to split.				
M67	Cnc	0	Contains 500+ stars mag 10 & fainter. One of the oldest clusters. Dist=2,350 ly.				
M94	CVn	0	Compact nearly face-on spiral galaxy. Dist=15 million ly.				
η Cassiopeiae	Cas	•	Yellow star mag 3.4 & orange star mag 7.5. Dist=19 ly. Orbit=480 years. Sep=12".				
5128	Cen	0	Bisected by a wide obscuring lane. Strong radio source. Dist=14 million ly.				
M51	CVn	0	Whirlpool Galaxy. First recognised to have spiral structure. Dist=25 million ly.				
M64	Com	0	Black-Eye Galaxy. Discovered by J.E. Bode in 1775 - "a small, nebulous star".				
Albireo	Cyg	•	Beautiful double star. Contrasting colours of orange and blue-green. Sep=34.4".				
61 Cygni	Cyg	•	Attractive double star. Mags 5.2 & 6.1 orange dwarfs. Dist=11.4 ly. Sep=28.4".				
3242	Hya		Ghost of Jupiter. Bright blue disk. Mag 11 central star. Dist=2,600 ly.				
M83	Hya	0	Classic face-on spiral. Discovered in 1752 by Lacaille. In attractive star field.				
γ Leonis	Leo	•	Superb pair of golden-yellow giant stars. Mags 2.2 & 3.5. Orbit=600 years. Sep=4.4".				
β Lyrae	Lyr	•	Eclipsing binary. Mag varies between 3.3 & 4.3 over 12.940 days. Fainter mag 7.2 blue star.				
M57	Lyr		Ring Nebula. Magnificent object. Smoke-ring shape. Dist=4,100 ly.				
M81	UMa	0	Beautiful spiral galaxy visible with binoculars. Easy to see in a telescope.				
M82	UMa	0	Close to M81 but much fainter and smaller.				
M87	Vir	0	Supergiant galaxy with supermassive black hole at its core. Dist=53.5 million ly.				

Sombrero Galaxy. Almost edge-on spiral galaxy. Protruding central core.

Superb pair of mag 3.5 yellow-white stars. Orbit=169 years. At their closest in 2005.

Dumbbell Nebula. Large, twin-lobed shape. Most spectacular planetary. Dist=975 ly.

The Evening Sky Map (ISSN 1839-7735) Copyright © 2000-2025 Kym Thalassoudis. All Rights Reserved.