

## AM, GM, Sequences, & Series Introduction

An ARITHMETIC SEQUENCE is one in which each consecutive terms share a common difference, i.e.

$$a, a + d, a + 2d, a + 3d, a + 4d, a + 5d, a + 6d, \dots \quad (\star)$$

The terms in an arithmetic sequence satisfy an interesting property.

1. Pick any term in the arithmetic sequence ( $\star$ ). How is it related to the term immediately before it and the term immediately after it?

It turns out, this has a name: for any two numbers  $x$  and  $y$ , the quantity  $\frac{x+y}{2}$  is the \_\_\_\_\_.

$$AM = \frac{x+y}{2}$$

More generally, for  $n$  numbers  $a_1, \dots, a_n$ , the arithmetic mean is

$$AM = \frac{a_1 + a_2 + \dots + a_n}{n}$$

2. (Competition Example) Let  $S = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ . If you randomly select 3 unique numbers from  $S$ , what is the probability the three numbers form an arithmetic progression?

---

A GEOMETRIC SEQUENCE is one in which each term is a constant multiple of the previous term, i.e.

$$a, ar, ar^2, ar^3, ar^4, ar^5, \dots ar^n, \dots \quad (\diamond)$$

The terms in a geometric sequence satisfy an interesting property.

3. Pick any term in the geometric sequence ( $\diamond$ ). How is it related to the *product* of the term immediately before it and the term immediately after it?

It turns out, this has a name: for any two numbers  $x$  and  $y$ , the quantity  $\sqrt{xy}$  is the \_\_\_\_\_.

$$GM = \sqrt{xy}$$

More generally, for  $n$  numbers  $a_1, \dots, a_n$ , the geometric mean is

$$GM = \sqrt[n]{a_1 a_2 \dots a_n}$$

4. (Competition Example) A geometric progression has 5 terms, two of which are known.

$$g_n = \{3, g_2, g_3, g_4, 27\}$$

Find  $g_4$ .

There is an interesting relationship between the arithmetic and geometric means of any given numbers. It is:

$$AM \geq GM$$

or

$$\frac{a_1 + a_2 + \cdots + a_n}{n} \geq \sqrt[n]{a_1 a_2 \cdots a_n}$$

and the two expressions will only be equal if \_\_\_\_\_. This theorem is of major use whenever we are trying to *minimize* or *maximize* a quantity.

5. (Competition Example) Let  $a, b, c$  be positive real numbers with  $abc = 1$ . Find the minimum possible of

$$S = (a + 1)(b + 1)(c + 1)$$

When we take a sequence and add all the terms, it is called a **SERIES**; arithmetic sequences become arithmetic series, and geometric sequences become geometric series. What primarily concerns us right now is finding values for these series quickly. Let's do some deriving.

### Arithmetic Series Formulas

1. (General Arithmetic Series)  $\sum_{k=0}^n (a + dk) =$

2. (Sum to  $n$ )  $\sum_{k=1}^n k =$

3. (Finite Geometric Series)  $\sum_{k=0}^n ar^k =$

4. (Infinite Geometric Series with  $|r| < 1$ ):  $\sum_{k=0}^{\infty} ar^k =$

Before some easier competition style problems, let's look at how the techniques from our derivations can solve a hard problem.

**Example:** Compute  $\sum_{n=1}^{\infty} \frac{n}{5^n}$ .

Finally, some more practice.

6. Compute  $1 - \frac{2}{3} + \frac{4}{9} - \frac{8}{27} + \dots$

7. Rebekah wanted to stump her teacher, so she whipped out a calculator and added up the integers from 12 to 88. Before she could finish saying "I just added from up 12 to 88, Mr. Smart Teacher Man. I bet you can't tell me...", the teacher said the correct answer. What was it?

8. (Competition Example) The numbers  $x, 18, y$ , in order, form an arithmetic progression. The numbers  $18, y, 90$ , in order, also form an arithmetic progression. What is  $x$ ?

9. (Competition Example) The terms of a geometric sequence  $a_n$  are all positive. Given that

$$a_2 + a_5 = 90 \quad \text{and} \quad a_3 a_7 = 6400,$$

find  $a_4$ .

**Homework:** Go to Alcumus ([artofproblemsolving.com](http://artofproblemsolving.com) → Resources (dropdown) → Alcumus), select "Algebra," and play any of

- Arithmetic Sequences
- Arithmetic Series
- Geometric Sequences
- Geometric Series

For more of a challenge, try the "Intermediate Algebra" focus and any of

- Arithmetic Sequences and Series
- Geometric Sequences and Series
- Telescoping Series (not covered today)
- General Sequences and Series
- Advanced Sequences and Series

## Solutions

An ARITHMETIC SEQUENCE is one in which each consecutive terms share a common difference, i.e.

$$a, a + d, a + 2d, a + 3d, a + 4d, a + 5d, a + 6d, \dots \quad (\star)$$

The terms in an arithmetic sequence satisfy an interesting property.

1. Pick any term in the arithmetic sequence ( $\star$ ). How is it related to the term immediately before it and the term immediately after it? **It is the average.**

It turns out, this has a name: for any two numbers  $x$  and  $y$ , the quantity  $\frac{x+y}{2}$  is the ARITHMETIC MEAN.

$$AM = \frac{x+y}{2}$$

More generally, for  $n$  numbers  $a_1, \dots, a_n$ , the arithmetic mean is

$$AM = \frac{a_1 + a_2 + \dots + a_n}{n}$$

2. Let the numbers selected be  $x < y < z$ . Then  $y = \frac{x+z}{2}$ , so  $x+z$  must be even and hence  $x$  and  $z$  must have the same parity. Therefore, out of the  $\binom{10}{3} = \frac{10(9)(8)}{3(2)(1)} = 120$  ways to choose 3 numbers, we can select any of the values of  $x+z$ , which will have a unique  $y$  and hence triplet  $(x, y, z)$ , and record how many possibilities there are for  $x$  and  $z$ :

$$x+z=4 \Rightarrow 1 \text{ options } ((1, 3))$$

$$x+z=6 \Rightarrow 2 \text{ options } ((1, 5), (2, 4))$$

$$x+z=8 \Rightarrow 3 \text{ options } ((1, 7), (2, 6), (3, 5))$$

$$x+z=10 \Rightarrow 4 \text{ options } ((1, 9), (2, 8), (3, 7), (4, 6))$$

By symmetry, there are also 10 options for the sums  $x+z$  of 12, 14, 16, and 18. Therefore, the probability of getting an arithmetic progression is  $\frac{20}{120} = \frac{1}{6}$ .

---

A GEOMETRIC SEQUENCE is one in which each term is a constant multiple of the previous term, i.e.

$$a, ar, ar^2, ar^3, ar^4, ar^5, \dots ar^n, \dots \quad (\diamond)$$

The terms in a geometric sequence satisfy an interesting property.

3. Pick any term in the geometric sequence ( $\diamond$ ). How is it related to the *product* of the term immediately before it and the term immediately after it? **It is the square root of the product.**

It turns out, this has a name: for any two numbers  $x$  and  $y$ , the quantity  $\sqrt{xy}$  is the GEOMETRIC MEAN.

$$GM = \sqrt{xy}$$

More generally, for  $n$  numbers  $a_1, \dots, a_n$ , the geometric mean is

$$GM = \sqrt[n]{a_1 a_2 \dots a_n}$$

4. We can quickly find  $g_3 = \sqrt{3(27)} = 9$  and  $g_4 = \sqrt{9(27)} = 9\sqrt{3}$ .

There is an interesting relationship between the arithmetic and geometric means of any given numbers. It is:

$$AM \geq GM$$

or

$$\frac{a_1 + a_2 + \dots + a_n}{n} \geq \sqrt[n]{a_1 a_2 \dots a_n}$$

and the two expressions will only be equal if \_\_\_\_\_ . This theorem is of major use whenever we are trying to *minimize* or *maximize* a quantity.

5. We use the AM-GM inequality on each of the terms  $a + 1, b + 1, c + 1$ .

$$\frac{a + 1}{2} \leq \sqrt{a(1)} \Rightarrow a + 1 \geq 2\sqrt{a}$$

$$\frac{b + 1}{2} \leq \sqrt{b(1)} \Rightarrow b + 1 \geq 2\sqrt{b}$$

$$\frac{c + 1}{2} \leq \sqrt{c(1)} \Rightarrow c + 1 \geq 2\sqrt{c}$$

Therefore,  $S \geq (2\sqrt{a})(2\sqrt{b})(2\sqrt{c}) = 8\sqrt{abc} = 8$ . The minimum value is 8.

When we take a sequence and add all the terms, it is called a SERIES; arithmetic sequences become arithmetic series, and geometric sequences become geometric series. What primarily concerns us right now is finding values for these series quickly. Let's do some deriving.

### Arithmetic Series Formulas

1. (General Arithmetic Series)  $\sum_{k=0}^n (a + dk) = \frac{(a + (a + dn)) \cdot \frac{n}{2}}{1} = \frac{(\text{first} + \text{last}) \cdot \frac{\# \text{ terms}}{2}}{1}$

2. (Sum to  $n$ )  $\sum_{k=1}^n k = \frac{n(n + 1)}{2}$

3. (Finite Geometric Series)  $\sum_{k=0}^n ar^k = a \cdot \frac{1 - r^{n+1}}{1 - r}$

4. (Infinite Geometric Series with  $|r| < 1$ ):  $\sum_{k=0}^{\infty} ar^k = a \cdot \frac{1}{1 - r}$

Before some easier competition style problems, let's look at how the techniques from our derivations can solve a hard problem.

**Example:** Let's call our sum  $S$ . Then

$$S = \frac{1}{5} + \frac{2}{25} + \frac{3}{125} + \frac{4}{625} + \dots$$

On first inspection, this series looks almost geometric, except for the pesky numerators, which won't stay constant. If we want to utilize the semi-geometric nature of the series, perhaps it would be useful to multiply it by  $\frac{1}{5}$ .

$$\frac{1}{5}S = \frac{1}{5} \left( \frac{1}{5} + \frac{2}{25} + \frac{3}{125} + \frac{4}{625} + \dots \right) = \frac{1}{25} + \frac{2}{125} + \frac{3}{625} + \dots$$

Aha! It looks just the original sum  $S$ , only the numerators have all been decreased by 1. This means, if we subtract the sums, we should get some useful simplification.

$$\begin{aligned} S - \frac{1}{5}S &= \left( \frac{1}{5} + \frac{2}{25} + \frac{3}{125} + \frac{4}{625} + \dots \right) - \left( \frac{1}{25} + \frac{2}{125} + \frac{3}{625} + \dots \right) \\ &= \frac{1}{5} + \frac{1}{25} + \frac{1}{125} + \frac{1}{625} + \dots \end{aligned}$$

We have our geometric series! We can now easily sum this and find  $S$ .

$$\begin{aligned} \frac{4}{5}S &= \frac{1}{5} \cdot \frac{1}{1 - \frac{1}{5}} \\ S &= \frac{5}{4} \cdot \frac{1}{5} \cdot \frac{5}{4} \\ S &= \frac{5}{16} \end{aligned}$$

Our series has a sum of  $\frac{5}{16}$ .

---

6. This is a geometric series with initial term 1 and ratio  $\frac{2}{3}$ , so it equals  $\frac{1}{1 - (-\frac{2}{3})} = \frac{3}{5}$ .

7.  $(12 + 88) \cdot \frac{(88 - 12 + 1)}{2} = \mathbf{3850}$

8. We have  $18 = \frac{x + y}{2}$  and  $y = \frac{18 + 90}{2}$ . Therefore,  $y = 54$  and  $x = -18$ .

9. The second equation, by the geometric mean, yields  $a_5 = \sqrt{6400} = 80$ . Plugging this into the first equation yields  $a_2 = 10$ . Then, the common ratio is clearly 2, so  $a_4 = 10(2)(2) = \mathbf{40}$ .